



ELSEVIER

Contents lists available at ScienceDirect

Journal of Functional Analysis

www.elsevier.com/locate/jfa



Diagonalization of the infinite q -boson system [☆]

J.F. van Diejen ^{*}, E. Emsiz

Facultad de Matemáticas, Pontificia Universidad Católica de Chile, Casilla 306, Correo 22, Santiago, Chile

ARTICLE INFO

Article history:

Received 17 October 2013

Accepted 23 January 2014

Available online 10 February 2014

Communicated by A. Borodin

Keywords:

 q -Bosons

Hall–Littlewood functions

 n -Particle scattering

ABSTRACT

We present a hierarchy of commuting operators in Fock space containing the q -boson Hamiltonian on \mathbb{Z} and show that the operators in question are simultaneously diagonalized by Hall–Littlewood functions and have absolutely continuous spectrum. As an application, the n -particle scattering operator is computed explicitly.

© 2014 Elsevier Inc. All rights reserved.

1. Introduction

The q -boson model constitutes a one-dimensional exactly solvable particle system in Fock space [1] based on the q -oscillator algebra [7, Ch. 5]. In the case of periodic boundary conditions (i.e. with particles hopping on the finite lattice \mathbb{Z}_m), the integrability, the spectrum, and the eigenfunctions of the Hamiltonian were studied by means of the algebraic Bethe Ansatz method [1]. Remarkably, these eigenfunctions turn out to be Hall–Littlewood functions at discrete spectral values [16] (cf. also Ref. [6] for an alternative construction of Hall–Littlewood functions in Fock space based on deformed vertex operator algebras, with applications in the study of KP τ -functions arising from

[☆] Work was supported in part by the *Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT)* Grants # 1130226 and # 11100315, and by the *Anillo ACT56 ‘Reticulados y Simetrías’* financed by the *Comisión Nacional de Investigación Científica y Tecnológica (CONICYT)*.

^{*} Corresponding author.

E-mail addresses: diejen@mat.puc.cl (J.F. van Diejen), eemsiz@mat.puc.cl (E. Emsiz).

generating functions of weighted plane partitions [5]). With the aid of explicit expressions for the commuting quantum integrals arising from an infinite-dimensional solution of the Yang–Baxter equation, it was recently demonstrated [8] that the eigenvalue problem for the q -boson system on \mathbb{Z}_m is in fact equivalent to that of an integrable discretization [3] of the celebrated delta Bose gas on the circle [9].

The current paper addresses the spectral problem and the integrability of the q -boson system on the *infinite* integer lattice \mathbb{Z} . Specifically, we demonstrate that the eigenfunctions of this infinite q -boson system are again given by Hall–Littlewood functions and provide explicit formulas for a complete hierarchy of operators commuting with the Hamiltonian; these formulas are natural infinite-dimensional analogues of the above-mentioned expressions in [8] for the finite periodic q -boson system on \mathbb{Z}_m . While in the case of the periodic lattice the use of the algebraic Bethe Ansatz method was instrumental for the diagonalization of the system, here in the infinite-dimensional setting our main tool consists of well-known Pieri formulas for the Hall–Littlewood functions [10]. The n -particle spectrum now turns out to be absolutely continuous rather than discrete but still remains bounded (as in the periodic case). The corresponding n -particle scattering operator relating the large-times asymptotics of the dynamics for $t \rightarrow -\infty$ and $t \rightarrow +\infty$ is computed explicitly as an application of Ruijsenaars’ general scattering results in Ref. [14].

The infinite q -boson model considered here turns out to be limiting case of a more general quantum particle system on \mathbb{Z} arising from a q -deformation of the totally asymmetric simple exclusion process (q -TASEP) [15,2]. The spectral problem for the latter q -boson system was solved very recently by means of a novel one-parameter inhomogeneous generalization of the Hall–Littlewood functions [2]. It is an open question whether the Fock space constructions of the present paper can be adapted so as to incorporate this more general q -boson model originating from the q -TASEP.

2. The infinite q -boson system

Given $n \geq 0$ integral, let $\mathcal{F}(A_n)$ be the space of complex functions $f : A_n \rightarrow \mathbb{C}$ on the fundamental domain

$$A_n := \{(\lambda_1, \dots, \lambda_n) \in \mathbb{Z}^n \mid \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n\} \quad (2.1)$$

of the integral lattice \mathbb{Z}^n modulo the action of the permutation group S_n , where $A_0 := \{\emptyset\}$ and $\mathcal{F}(A_0) := \mathbb{C}$ by convention. We will refer to the infinite direct sum

$$\mathcal{F} := \bigoplus_{n \geq 0} \mathcal{F}(A_n) \quad (2.2)$$

built of all *finite* linear combinations of (arbitrary) functions $f_n \in \mathcal{F}(A_n)$, $n = 0, 1, 2, \dots$ as the algebraic Fock space.

For $\lambda \in A_n$ and $l \in \mathbb{Z}$, let the multiplicity $m_l(\lambda)$ count the number of components λ_j , $1 \leq j \leq n$ such that $\lambda_j = l$. We write $\beta_l^* \lambda$ for the point of A_{n+1} obtained from λ by inserting an additional component with value l and—assuming $m_l(\lambda) > 0$ —we write $\beta_l \lambda \in A_{n-1}$ for the result of the inverse operation that deletes a component with value l from λ . Upon defining the following actions on $f \in \mathcal{F}(A_n)$:

$$\begin{aligned}
 (\beta_l f)(\lambda) &:= \begin{cases} f(\beta_l^* \lambda) & \text{if } n > 0 \quad (\lambda \in A_{n-1}), \\ 0 & \text{if } n = 0, \end{cases} \\
 (\beta_l^* f)(\lambda) &:= \begin{cases} [m_l(\lambda)]f(\beta_l \lambda) & \text{if } m_l(\lambda) > 0 \\ 0 & \text{otherwise} \end{cases} \quad (\lambda \in A_{n+1}), \\
 (N_l f)(\lambda) &:= q^{m_l(\lambda)} f(\lambda) \quad (\lambda \in A_n),
 \end{aligned} \tag{2.3}$$

where $0 < q < 1$ and

$$[m] := \frac{1 - q^m}{1 - q} = 1 + q + \dots + q^{m-1}$$

for $m = 0, 1, 2, \dots$, it is readily verified that one ends up with a representation of the q -boson field algebra on \mathcal{F} :

$$[\beta_l, \beta_k] = [\beta_l^*, \beta_k^*] = [N_l, N_k] = [N_l, \beta_k] = [N_l, \beta_k^*] = [\beta_l, \beta_k^*] = 0 \tag{2.4a}$$

for $l \neq k$, and

$$N_l \beta_l^* = q \beta_l^* N_l, \quad \beta_l N_l = q N_l \beta_l, \quad [\beta_l, \beta_l^*] = N_l, \quad \beta_l \beta_l^* - q \beta_l^* \beta_l = 1. \tag{2.4b}$$

Here the brackets refer to the (ordinary) commutator product. By construction β_l, β_l^* and N_l map $\mathcal{F}(A_n)$ into $\mathcal{F}(A_{n-1}), \mathcal{F}(A_{n+1})$ and $\mathcal{F}(A_n)$, respectively (with the convention that $\mathcal{F}(A_{-1}) = \{0\}$).

The Hamiltonian of the q -boson system

$$H_q = \sum_{l \in \mathbb{Z}} (a_l + a_l^*) \tag{2.5a}$$

is built of hopping operators

$$a_l := \beta_{l+1}^* \beta_l \quad \text{and} \quad a_l^* := \beta_{l+1} \beta_l^* \tag{2.5b}$$

for which the n -particle subspace $\mathcal{F}(A_n)$ is stable. These hopping operators represent the plactic subalgebra of the q -boson field algebra [8, Sec. 3.4]:

$$a_l a_k = a_k a_l \tag{2.6a}$$

for $|l - k| > 1$ (nonlocal commutativity) and

$$\begin{aligned}
 a_{l+1}a_l^2 + qa_l^2a_{l+1} &= (1+q)a_la_{l+1}a_l, \\
 a_{l+1}^2a_l + qa_la_{l+1}^2 &= (1+q)a_{l+1}a_la_{l+1}
 \end{aligned}
 \tag{2.6b}$$

(quantum Knuth relations), with analogous relations (involving reversely ordered products) for a_l^* , $l \in \mathbb{Z}$. The q -boson Hamiltonian H_q (2.5a), (2.5b) constitutes a well-defined operator on \mathcal{F} as for any $f \in \mathcal{F}(\Lambda_n)$ and $\lambda \in \Lambda_n$ the infinite sum $(H_q f)(\lambda)$ contains only a finite number of nonvanishing terms.

To facilitate the comparison with previous literature on the q -boson system [1,16,8], let us denote the characteristic function in $\mathcal{F}(\Lambda_n)$ supported on $\lambda \in \Lambda_n$ by $|\lambda\rangle$. Then one has that

$$\beta_l|\lambda\rangle = \begin{cases} |\beta_l\lambda\rangle & \text{if } m_l(\lambda) > 0, \\ 0 & \text{otherwise,} \end{cases} \quad \beta_l^*|\lambda\rangle = [m_l(\lambda) + 1]|\beta_l^*\lambda\rangle, \quad N_l|\lambda\rangle = q^{m_l(\lambda)}|\lambda\rangle.$$

In the standard physical interpretation the state $|\lambda\rangle$ encodes a configuration of n particles on \mathbb{Z} — q -bosons—with $m_l(\lambda)$ particles occupying the site $l \in \mathbb{Z}$. The operators β_l^* and β_l play the role of particle creation and annihilation operators and N_l counts the number of particles at the site l (as a power of q). The hopping operators a_l and a_l^* move a particle from l to $l + 1$ and vice versa.

3. Integrability

To any partition $\eta = (\eta_1, \dots, \eta_p)$ with $\eta_1 \geq \eta_2 \geq \dots \geq \eta_p \geq 1$, we associate the following hopping operators on \mathcal{F} :

$$\begin{aligned}
 m_\eta(a) &:= \sum'_{\sigma \in S_p} \sum_{l_1 < l_2 < \dots < l_p} a_{l_1}^{\eta_{\sigma_1}} \dots a_{l_p}^{\eta_{\sigma_p}}, \\
 m_\eta(a^*) &:= \sum'_{\sigma \in S_p} \sum_{l_1 > l_2 > \dots > l_p} (a_{l_1}^*)^{\eta_{\sigma_1}} \dots (a_{l_p}^*)^{\eta_{\sigma_p}}.
 \end{aligned}
 \tag{3.1}$$

Here the (infinite) inner summations are over all strictly monotonous p -tuples (l_1, \dots, l_p) of indices in \mathbb{Z} ; the primes attached to the (finite) outer summations indicate that these are meant over the orbit of all distinct compositions $(\eta_{\sigma_1}, \dots, \eta_{\sigma_p})$ obtained by reordering the parts of η via permutations

$$\sigma = \begin{pmatrix} 1 & 2 & \dots & p \\ \sigma_1 & \sigma_2 & \dots & \sigma_p \end{pmatrix}$$

belonging to the symmetric group S_p . Notice that for given $f \in \mathcal{F}(\Lambda_n)$ and $\lambda \in \Lambda_n$, the infinite sums $(m_\eta(a)f)(\lambda)$ and $(m_\eta(a^*)f)(\lambda)$ contain only a finite number of nonzero terms, so these operators are again well-defined on \mathcal{F} . For $r \in \mathbb{N}$ we now set

$$H_r := \sum_{|\eta|=r} \frac{m_\eta(a)}{[\eta]!} \quad \text{and} \quad H_r^* := \sum_{|\eta|=r} \frac{m_\eta(a^*)}{[\eta]!}, \tag{3.2}$$

where $[\eta]! = [(\eta_1, \dots, \eta_p)]! := [\eta_1]! \cdots [\eta_p]!$ and $[m]! := [m][m-1] \cdots [1]$ for $m = 0, 1, 2, \dots$ (with the convention that $[0]! = 1$), and $|\eta| := \eta_1 + \dots + \eta_p$ (so the (finite) summation in Eq. (3.2) is over all partitions of r). The q -boson Hamiltonian (2.5a), (2.5b) becomes in terms of these operators:

$$H_q = H_1 + H_1^*. \tag{3.3}$$

Our main result is the following explicit formula for the action of H_r and H_r^* in the n -particle subspace $\mathcal{F}(\Lambda_n)$, which will be proven shortly in the next section.

Theorem 3.1 (*Explicit action of $H_r^{(*)}$ in $\mathcal{F}(\Lambda_n)$*). *For any $f \in \mathcal{F}(\Lambda_n)$ and $\lambda \in \Lambda_n$, one has that*

$$\begin{aligned} (H_r f)(\lambda) &= \sum_{\substack{J \subset \{1, 2, \dots, n\}, |J|=r \\ \lambda - e_J \in \Lambda_n}} V_{\lambda, J^c} f(\lambda - e_J), \\ (H_r^* f)(\lambda) &= \sum_{\substack{J \subset \{1, 2, \dots, n\}, |J|=r \\ \lambda + e_J \in \Lambda_n}} V_{\lambda, J} f(\lambda + e_J), \end{aligned} \tag{3.4a}$$

where $|J|$ denotes the cardinality of $J \subset \{1, \dots, n\}$, $J^c := \{1, \dots, n\} \setminus J$, $e_J := \sum_{j \in J} e_j$ (with e_1, \dots, e_n referring to the standard unit basis of \mathbb{Z}^n) and

$$V_{\lambda, J} := \prod_{\substack{1 \leq j < k \leq n \\ j \in J, k \in J^c \\ \lambda_j = \lambda_k}} \frac{1 - q^{k-j+1}}{1 - q^{k-j}}. \tag{3.4b}$$

In particular, for $r > n$ the n -particle subspace $\mathcal{F}(\Lambda_n)$ belongs to the kernel of the operators H_r and H_r^* .

In [4, App. C] the discrete difference operators H_r on the RHS of Eqs. (3.4a), (3.4b) were obtained (up to a trivial similarity transformation and replacing q by q^2) as n algebraically independent commuting central elements arising from a difference-reflection representation of the affine Hecke algebra associated with $GL(n; \mathbb{C})$. Since H_n acts on $f \in \mathcal{F}(\Lambda_n)$ simply as an overall translational symmetry: $(H_n f)(\lambda) = f(\lambda - (e_1 + \dots + e_n))$, it is immediate from the above formulas that the discrete difference operators in $\mathcal{F}(\Lambda_n)$ corresponding to H_r^* can be written in turn as $H_{n-r} H_n^{-1}$ (with the convention that $H_0 := 1$). The upshot is that all H_r and H_r^* (3.2) commute as operators on \mathcal{F} , which proves the integrability of the infinite q -boson system.

Corollary 3.2 (Integrability). All operators H_r and H_r^* in Eq. (3.2) mutually commute on \mathcal{F} :

$$[H_r, H_{r'}] = 0, \quad [H_r^*, H_{r'}^*] = 0, \quad [H_r, H_{r'}^*] = 0 \quad (\forall r, r' \in \mathbb{N}), \quad (3.5)$$

and they restrict to n algebraically independent operators on the invariant n -particle subspace $\mathcal{F}(A_n)$.

A quasi-periodic counterpart of the formula in Theorem 3.1 for the finite q -boson system on \mathbb{Z}_m can be found in Ref. [8] (see Prp. 3.11 and Prp. 6.1). For Dirichlet type boundary conditions corresponding to the case of a vanishing quasi-periodicity parameter, one arrives—in the limit when the lattice size parameter m tends to infinity—at an analogue of the commutativity in Corollary 3.2 for the q -boson system on the (semi-)infinite lattice \mathbb{N} as a consequence of [8, Cor. 3.3] (cf. also [3, Thm. 5.3]). In principle, the commutativity in Corollary 3.2 for the q -boson system on \mathbb{Z} could also be recovered along these lines upon centering the finite lattice around the origin before performing the infinite size limit. Alternatively, the commutativity in question can also be viewed as a degeneration of the commutativity of the discrete Macdonald–Ruijsenaars operators [13], [10, Sec. VI.6] via a limit transition (that takes Macdonald symmetric functions to Hall–Littlewood symmetric functions).

It is straightforward from Theorem 3.1 and Eq. (3.3) that the action of the q -boson Hamiltonian H_q (2.5a), (2.5b) in the n -particle subspace $\mathcal{F}(A_n)$ is given by

$$(H_q f)(\lambda) = \sum_{\substack{1 \leq j \leq n, \epsilon = \pm 1 \\ \lambda + \epsilon e_j \in A_n}} [m_{\lambda_j}(\lambda)] f(\lambda + \epsilon e_j) \quad (3.6)$$

($f \in \mathcal{F}(A_n)$, $\lambda \in A_n$).

4. Proof of the main Theorem 3.1

We will determine the action of H_r^* on $f \in \mathcal{F}_n$ by direct computation in three steps; the calculation of $H_r f$ is completely analogous so its details will be suppressed (but cf. Remark 6.3 below for an alternative shortcut yielding the action of H_r from that of H_r^* via adjointness).

4.1. In the first step $V_{\lambda, J}$ (3.4b) (with $\lambda, \lambda + e_J \in A_n$) is rewritten in q -binomial form by means of well-known product formulas for the Poincaré polynomial of the symmetric group:

$$S_n(q) := \sum_{\sigma \in S_n} q^{\ell(\sigma)} = \prod_{1 \leq j < k \leq n} \frac{1 - q^{1+k-j}}{1 - q^{k-j}} = \prod_{1 \leq j \leq n} \frac{1 - q^j}{1 - q} = [n]! \quad (4.1a)$$

(where $\ell(\sigma)$ denotes the *length* of σ). Since the stabilizer subgroup $S_{n,\lambda} := \{\sigma \in S_n \mid \sigma\lambda = \lambda\}$ is isomorphic to the direct product $\prod_{l \in \mathbb{Z}} S_{m_l(\lambda)}$, the corresponding Poincaré polynomial factorizes in turn as

$$S_{n,\lambda}(q) = \sum_{\substack{\sigma \in S_n \\ \sigma\lambda = \lambda}} q^{\ell(\sigma)} = \prod_{\substack{1 \leq j < k \leq n \\ \lambda_j = \lambda_k}} \frac{1 - q^{1+k-j}}{1 - q^{k-j}} = \prod_{l \in \mathbb{Z}} [m_l(\lambda)]! \tag{4.1b}$$

and similarly $S_{n,\lambda} \cap S_{n,\lambda+e_J} \cong \prod_{l \in \mathbb{Z}} (S_{m_{l,J}(\lambda)} \times S_{m_{l,J^c}(\lambda)})$ so

$$\begin{aligned} (S_{n,\lambda} \cap S_{n,\lambda+e_J})(q) &= \prod_{\substack{j,k \in J \\ j < k, \lambda_j = \lambda_k}} \frac{1 - q^{1+k-j}}{1 - q^{k-j}} \prod_{\substack{j,k \in J^c \\ j < k, \lambda_j = \lambda_k}} \frac{1 - q^{1+k-j}}{1 - q^{k-j}} \\ &= \prod_{l \in \mathbb{Z}} [m_{l,J}(\lambda)]! [m_{l,J^c}(\lambda)]!, \end{aligned} \tag{4.1c}$$

where $m_{l,J}(\lambda)$ denotes the number of components λ_j , $j \in J$ such that $\lambda_j = l$ (i.e. $m_{l,J}(\lambda) + m_{l,J^c}(\lambda) = m_l(\lambda)$). Division of Eqs. (4.1b) and (4.1c) now reveals that

$$V_{\lambda,J} = V_{\lambda,J} V_{\lambda,J^c} = \frac{S_{n,\lambda}(q)}{(S_{n,\lambda} \cap S_{n,\lambda+e_J})(q)} = \prod_{l \in \mathbb{Z}} \left[\frac{m_l(\lambda)}{m_{l,J}(\lambda)} \right] \tag{4.2}$$

where $\left[\frac{m}{k} \right] := \frac{[m]!}{[k]![m-k]!}$ for $m \geq k \geq 0$. (Notice in this connection that here $V_{\lambda,J^c} = 1$, because the product in question is *empty* as consequence of the assumption that $\lambda + e_J$ belongs to A_n .)

4.2. It is clear by induction on $m \geq 1$ that for any $f \in \mathcal{F}_n$ and $\lambda \in A_n$:

$$((a_l^*)^m f)(\lambda) = \begin{cases} [m]! \left[\frac{m_l(\lambda)}{m} \right] f(a_l^m \lambda) & \text{if } m \leq m_l(\lambda), \\ 0 & \text{if } m > m_l(\lambda), \end{cases} \tag{4.3}$$

where in the former case $a_l^m \lambda = \lambda + e_d + e_{d+1} + \dots + e_{d+m-1}$ —with $d = d(\lambda, l) := \min\{j \mid \lambda_j = l\}$ —belongs to A_n (because of the condition that $m \leq m_l(\lambda)$). By iterating the formula in Eq. (4.3) it readily follows that—for $l_1 > l_2 > \dots > l_p$ and a composition (m_1, m_2, \dots, m_p) obtained by reordering the parts of the partition $\eta = (\eta_1, \eta_2, \dots, \eta_p)$ (with $\eta_p \geq 1$)—the action of the corresponding monomial in $m_\eta(a^*)$ (3.1) is given by

$$\begin{aligned} ((a_{l_1}^*)^{m_1} \dots (a_{l_p}^*)^{m_p} f)(\lambda) &= [\eta]! \left[\frac{m_{l_1}(\lambda)}{m_1} \right] \dots \left[\frac{m_{l_p}(\lambda)}{m_p} \right] f(a_{l_p}^{m_p} \dots a_{l_1}^{m_1} \lambda) \\ &= [\eta]! \prod_{l \in \mathbb{Z}} \left[\frac{m_l(\lambda)}{m_{l,J}(\lambda)} \right] f(\lambda + e_J) \end{aligned} \tag{4.4}$$

provided $m_k \leq m_{l_k}(\lambda)$ for $k = 1, \dots, p$, and equal to zero otherwise. Here

$$J = \{d_k, d_k + 1, \dots, d_k + m_k - 1 \mid k = 1, \dots, p\} \tag{4.5}$$

with $d_k := d(\lambda, l_k) = \min\{j \mid \lambda_j = l_k\}$. The condition that $m_k \leq m_{l_k}(\lambda)$ for $k = 1, \dots, p$ guarantees that J is a subset of $\{1, \dots, n\}$ of cardinality $|J| = m_1 + \dots + m_p = |\eta|$ and that $\lambda + e_J = a_{l_p}^{m_p} \cdots a_{l_1}^{m_1} \lambda \in A_n$.

4.3. From Steps 4.1 and 4.2 one learns that for $r \leq n$ the action $(H_r^* f)(\lambda)$ is built of a sum of terms of the form $V_{\lambda, J} f(\lambda + e_J)$ (cf. Eqs. (4.2) and (4.4)), with $J \subset \{1, \dots, n\}$ satisfying that $|J| = r$ and $\lambda + e_J \in A_n$. For $r > n$ on the other hand $(H_r^* f)(\lambda)$ vanishes (since then for all monomial terms $m_1 + \dots + m_p = |\eta| = r > n \geq m_{l_1}(\lambda) + \dots + m_{l_p}(\lambda)$).

To complete the proof of the explicit formula for $H_r^* f$ in Theorem 3.1, it only remains to infer that in the former situation all terms on the RHS actually do occur and with multiplicity 1. Indeed, this is clear from the observation that given $\lambda \in A_n$ and $J \subset \{1, \dots, n\}$ such that $\lambda + e_J \in A_n$, the corresponding $l_1 > l_2 > \dots > l_p$ and m_1, m_2, \dots, m_p for which

$$([m_1]! \cdots [m_p]!)^{-1} ((a_{l_1}^*)^{m_1} \cdots (a_{l_p}^*)^{m_p} f)(\lambda) = V_{\lambda, J} f(\lambda + e_J)$$

are uniquely retrieved by ordering the elements of the set $\{\lambda_j \mid j \in J\} = \{l_1, \dots, l_p\}$ and picking $m_k = m_{l_k, J}(\lambda)$, $k = 1, \dots, p$.

5. Diagonalization

For $\lambda \in A_n$ and a spectral parameter $\xi = (\xi_1, \xi_2, \dots, \xi_n)$ taken from the open fundamental alcove

$$A := \{\xi \in \mathbb{R}^n \mid \pi > \xi_1 > \xi_2 > \dots > \xi_n > -\pi\}, \tag{5.1}$$

let us define the n -variable Hall–Littlewood function as [10, Ch. III]

$$\phi_\xi(\lambda) := \sum_{\sigma \in S_n} C(\xi_\sigma) e^{i\lambda \cdot \xi_\sigma}, \tag{5.2a}$$

with $\xi_\sigma := (\xi_{\sigma_1}, \xi_{\sigma_2}, \dots, \xi_{\sigma_n})$ and

$$C(\xi) := \prod_{1 \leq j < k \leq n} \frac{1 - qe^{i(\xi_k - \xi_j)}}{1 - e^{i(\xi_k - \xi_j)}}. \tag{5.2b}$$

It is immediate from the explicit action in Theorem 3.1 and the Pieri formulas for the Hall–Littlewood functions [10, Sec. III.3] that the action of the commuting operators H_r and H_r^* (3.2) in the n -particle subspace $\mathcal{F}(A_n)$ is diagonal on ϕ_ξ (5.2a), (5.2b).

Corollary 5.1 (*Diagonalization*). *For any spectral value ξ in the fundamental alcove A (5.1), the n -variable Hall–Littlewood function ϕ_ξ (5.2a), (5.2b) constitutes a joint eigenfunction for the commuting operators H_r and H_r^* (3.2) in $\mathcal{F}(A_n)$:*

$$H_r \phi_\xi = e_r(e^{-i\xi}) \phi_\xi \quad \text{and} \quad H_r^* \phi_\xi = e_r(e^{i\xi}) \phi_\xi \quad (r = 1, \dots, n), \tag{5.3a}$$

with $e_r(e^{-i\xi}) := e_r(e^{-i\xi_1}, \dots, e^{-i\xi_n})$ and $e_r(e^{i\xi}) := e_r(e^{i\xi_1}, \dots, e^{i\xi_n})$, where e_r refers to the r th elementary symmetric function

$$e_r(x_1, \dots, x_n) := \sum_{1 \leq j_1 < j_2 < \dots < j_r \leq n} x_{j_1} x_{j_2} \cdots x_{j_r}. \tag{5.3b}$$

Proof. By Theorem 3.1, the eigenvalue equations in Eqs. (5.3a), (5.3b) become explicitly:

$$\begin{aligned} e_r(e^{-i\xi}) \phi_\xi(\lambda) &= \sum_{\substack{J \subset \{1, 2, \dots, n\}, |J|=r \\ \lambda - e_J \in A_n}} V_{\lambda, J^c} \phi_\xi(\lambda - e_J), \\ e_r(e^{i\xi}) \phi_\xi(\lambda) &= \sum_{\substack{J \subset \{1, 2, \dots, n\}, |J|=r \\ \lambda + e_J \in A_n}} V_{\lambda, J} \phi_\xi(\lambda + e_J), \end{aligned}$$

respectively. Both formulas boil down to well-known Pieri identities for the Hall–Littlewood functions [10, Sec. III.3]. In the form stated above the second identity can e.g. be directly retrieved from [4, Eq. (C.11)] and the first identity follows for $r < n$ from the second upon dividing by $e_n(e^{i\xi})$ and replacing r by $n - r$, whereas for $r = n$ both identities are equivalent and reduce to the elementary translational quasi-periodicity $\phi_\xi(\lambda + e_1 + \dots + e_n) = e^{i\xi_1 + \dots + i\xi_n} \phi_\xi(\lambda)$ (which is manifest from Eq. (5.2a)). \square

It follows in particular that the Hall–Littlewood function ϕ_ξ (5.2a), (5.2b) is an eigenfunction of the q -boson Hamiltonian H_q (2.5a), (2.5b) in the n -particle subspace $\mathcal{F}(A_n)$:

$$H_q \phi_\xi = \varepsilon(\xi) \phi_\xi \quad \text{with} \quad \varepsilon(\xi) := 2 \sum_{j=1}^n \cos(\xi_j) \tag{5.4}$$

(cf. Eq. (3.3)).

6. Spectral analysis

To address the completeness of the above eigenfunctions for the infinite q -boson system, we pass from our algebraic Fock space \mathcal{F} (2.2) to a full-fledged Fock space

$$\mathcal{H} := \bigoplus_{n \geq 0} \ell^2(A_n, \delta_n), \tag{6.1}$$

which is built of all series $\sum_{n \geq 0} f_n$, with $f_n \in \ell^2(\Lambda_n, \delta_n)$ such that $\sum_{n \geq 0} \langle f_n, f_n \rangle_n < \infty$. Here the n -particle Hilbert space $\ell^2(\Lambda_n, \delta_n) = \mathcal{H} \cap \mathcal{F}(\Lambda_n)$ consists of the functions $f \in \mathcal{F}(\Lambda_n)$ such that $\langle f, f \rangle_n < \infty$, where

$$\langle f, g \rangle_n := \sum_{\lambda \in \Lambda_n} f(\lambda) \overline{g(\lambda)} \delta_n(\lambda) \quad (f, g \in \ell^2(\Lambda_n, \delta_n)) \tag{6.2a}$$

with

$$\delta_n(\lambda) := 1/S_{n,\lambda}(q) = 1/\prod_{l \in \mathbb{Z}} [m_l(\lambda)]! \tag{6.2b}$$

(cf. Eq. (4.1b)).

The representation of the q -boson algebra in Eq. (2.3) readily extends from the dense domain

$$\mathcal{D} := \mathcal{H} \cap \mathcal{F} \tag{6.3}$$

consisting of the *finite* linear combinations $\sum_{n \geq 0} c_n f_n$ —with $c_n \in \mathbb{C}$ and $f_n \in \ell^2(\Lambda_n, \delta_n)$ —to a bounded representation on the Fock space \mathcal{H} (6.1). Indeed, it is immediate from the definitions that for any $f \in \ell^2(\Lambda_n, \delta_n)$:

$$\begin{aligned} \langle \beta_l f, \beta_l f \rangle_{n-1} &\leq (1-q)^{-1} \langle f, f \rangle_n, \\ \langle \beta_l^* f, \beta_l^* f \rangle_{n+1} &\leq (1-q)^{-1} \langle f, f \rangle_n, \\ \langle N_l f, N_l f \rangle_n &\leq \langle f, f \rangle_n \end{aligned} \tag{6.4}$$

(where one exploits that $\delta_{n+1, \beta_l^* \lambda}(q) = \delta_{n,\lambda}(q)/[m_l(\lambda) + 1]$ for all $\lambda \in \Lambda_n$ and that $[m] \leq 1/(1-q)$ for all $m = 0, 1, 2, \dots$). The representation at issue moreover preserves the $*$ -structure:

$$\langle \beta_l^* f, g \rangle_{n+1} = \langle f, \beta_l g \rangle_n \quad \text{and} \quad \langle N_l f, g \rangle_n = \langle f, N_l g \rangle_n \tag{6.5}$$

(for all $f \in \ell^2(\Lambda_n, \delta_n)$ with $g \in \ell^2(\Lambda_{n+1}, \delta_{n+1})$ and $g \in \ell^2(\Lambda_n, \delta_n)$, respectively).

The completeness of the eigenfunctions in Corollary 5.1 is now obvious from the well-known fact that the Hall–Littlewood functions $\phi_\xi(\lambda)$, $\lambda \in \Lambda_n$ form an orthogonal basis for the Hilbert space $L^2(A, \Delta d\xi)$ with inner product

$$\langle \hat{f}, \hat{g} \rangle_\Delta = \frac{1}{(2\pi)^n} \int_A \hat{f}(\xi) \overline{\hat{g}(\xi)} \Delta(\xi) d\xi, \quad \text{where } \Delta(\xi) := \frac{1}{|C(\xi)|^2} \tag{6.6}$$

with $C(\xi)$ taken from Eq. (5.2b). More specifically, for any $\lambda, \mu \in \Lambda_n$ one has that [11, §10]

$$\langle \phi(\lambda), \phi(\mu) \rangle_\Delta = \begin{cases} 1/\delta_n(\lambda) & \text{if } \lambda = \mu, \\ 0 & \text{otherwise.} \end{cases} \tag{6.7}$$

The corresponding Fourier transform $\mathbf{F}_q : \ell^2(\Lambda_n, \delta_n) \rightarrow L^2(A, \Delta d\xi)$ defined by

$$(\mathbf{F}_q f)(\xi) := \langle f, \phi_\xi \rangle_n = \sum_{\lambda \in \Lambda_n} f(\lambda) \overline{\phi_\xi(\lambda)} \delta_n(\lambda) \tag{6.8a}$$

($f \in \ell^2(\Lambda_n, \delta_n)$) thus determines a Hilbert space isomorphism with the inversion formula given by

$$(\mathbf{F}_q^{-1} \hat{f})(\lambda) = \langle \hat{f}, \overline{\phi(\lambda)} \rangle_\Delta = \frac{1}{(2\pi)^n} \int_A \hat{f}(\xi) \phi_\xi(\lambda) \Delta(\xi) d\xi \tag{6.8b}$$

($\hat{f} \in L^2(A, \Delta d\xi)$).

By [Corollary 5.1](#), this means that in the n -particle subspace $\ell^2(\Lambda_n, \delta_n)$ the higher commuting q -boson Hamiltonians

$$H_{q,r} := H_r + H_r^*, \quad r = 1, \dots, n, \tag{6.9}$$

are unitarily equivalent to bounded self-adjoint multiplication operators $\hat{E}_1, \dots, \hat{E}_n$ on $L^2(A, \Delta d\xi)$ of the form

$$(\hat{E}_r \hat{f})(\xi) := \varepsilon_r(\xi) \hat{f}(\xi) \tag{6.10a}$$

with

$$\varepsilon_r(\xi) := 2 \sum_{1 \leq j_1 < j_2 < \dots < j_r \leq n} \cos(\xi_{j_1} + \xi_{j_2} + \dots + \xi_{j_r}), \tag{6.10b}$$

viz.

$$H_{q,r} = \mathbf{F}_q^{-1} \circ \hat{E}_r \circ \mathbf{F}_q, \quad r = 1, \dots, n, \tag{6.11}$$

on $\ell^2(\Lambda_n, \delta_n)$.

Theorem 6.1 (*Spectral decomposition in $\ell^2(\Lambda_n, \delta_n)$*). *The higher q -boson Hamiltonians $H_{q,1}, \dots, H_{q,n}$ (6.9) constitute n independent commuting bounded self-adjoint operators on $\ell^2(\Lambda_n, \delta_n)$ with purely absolutely continuous spectrum. The spectral decomposition of these Hamiltonians in the n -particle subspace $\ell^2(\Lambda_n, \delta_n)$ is given explicitly by Eq. (6.11).*

As a consequence, the infinite q -boson hierarchy

$$H_{q,r} = H_r + H_r^*, \quad r \in \mathbb{N} \tag{6.12}$$

consists in turn of (commuting) symmetric operators in Fock space on the dense domain \mathcal{D} (6.3). It is not difficult to infer that these operators are actually essentially

self-adjoint and unbounded in the Fock space \mathcal{H} (6.1). Indeed, it is clear from the first part of Theorem 6.1 that for $z \in \mathbb{C} \setminus \mathbb{R}$ the operator $(H_{q,r} - z)$ maps the n -particle subspace $\ell^2(\Lambda_n, \delta_n)$ onto itself, whence the range $(H_{q,r} - z)\mathcal{D}$ is dense in \mathcal{H} . In other words, the deficiency indices of $H_{q,r}$ on the domain $\mathcal{D} \subset \mathcal{H}$ vanish and the claimed essential self-adjointness follows. The unboundedness of $H_{q,r}$ in \mathcal{H} is seen in turn from the last part of Theorem 6.1 and the observation that the n -particle spectrum does not remain bounded as n grows large: $\lim_{n \rightarrow \infty} \sup_{\xi \in A} |\varepsilon_r(\xi)| = \infty$. The upshot is that we can adapt the integrability result in Corollary 3.2 to the present setting as follows.

Theorem 6.2 (Integrability). *The higher Hamiltonians $H_{q,r}$ (6.12) of the infinite q -boson hierarchy on \mathcal{D} (6.3) extend uniquely to independent unbounded self-adjoint operators in the Fock space \mathcal{H} (6.1) with commuting resolvents $(H_{q,r} - z)^{-1}$ (with $z \in \mathbb{C} \setminus \mathbb{R}$ and $r \in \mathbb{N}$).*

Remark 6.3. It is a priori clear already from the definition of H_r and H_r^* (3.2) in terms of hopping operators—without need to resort to the explicit formula in Theorem 3.1—that for given $\lambda \in \Lambda_n$ and any $f \in \mathcal{F}(\Lambda_n)$ the values of $(H_r f)(\lambda)$ and $(H_r^* f)(\lambda)$ involve only evaluations of f at a finite number of points in Λ_n . Hence, for f in the subspace $C_0(\Lambda_n) \subset \mathcal{F}(\Lambda_n) \cap \ell^2(\Lambda_n, \delta_n)$ of functions with finite support in Λ_n the infinite sums comprising $H_r f$ and $H_r^* f$ contain only a finite number of nonvanishing monomial terms. The symmetry

$$\langle H_r f, g \rangle_n = \langle f, H_r^* g \rangle_n \quad (\forall f, g \in C_0(\Lambda_n)) \tag{6.13}$$

then follows from the first relation in Eq. (6.5) (without invoking the spectral decomposition in Eq. (6.11)). One can thus determine the action of H_r on $C_0(\Lambda_n)$ from the action of H_r^* (and vice versa) by computing the adjoint with respect to the inner product $\langle \cdot, \cdot \rangle_n$:

$$\begin{aligned} \langle H_r f, g \rangle_n &= \langle f, H_r^* g \rangle_n = \sum_{\lambda \in \Lambda_n} \delta_n(\lambda) f(\lambda) \overline{(H_r^* g)(\lambda)} \\ &= \sum_{\lambda \in \Lambda_n} \delta_n(\lambda) f(\lambda) \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \lambda + e_J \in \Lambda_n}} V_{\lambda, J} \overline{g(\lambda + e_J)} \\ &= \sum_{\lambda \in \Lambda_n} \delta_n(\lambda) \overline{g(\lambda)} \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \lambda - e_J \in \Lambda_n}} V_{\lambda, J} f(\lambda - e_J), \end{aligned}$$

where it was used in the last step that $\delta_n(\lambda - e_J) V_{\lambda - e_J, J} = \delta_n(\lambda) V_{\lambda, J^c}$ (for $\lambda \in \Lambda_n$ such that $\lambda - e_J \in \Lambda_n$). Since the actions of H_r and H_r^* on $\mathcal{F}(\Lambda_n)$ are determined completely by their restrictions to the subspace $C_0(\Lambda_n)$ (by the opening statement of this remark), the above computation shows that both formulas in Theorem 3.1 follow from each other (so it indeed suffices in the proof of Theorem 3.1 to verify only one of these two cases directly).

7. *n*-Particle scattering

For $q \rightarrow 0$, the Hall–Littlewood functions (5.2a), (5.2b) reduce to Schur functions [10, Ch. I]; the q -boson system degenerates in this limit to a system of impenetrable bosons known as the *Phase Model* [1,16,8]. We end up by computing the scattering operator that compares the large-time asymptotics of the n -particle dynamics of the q -boson system with that of the phase model.

In Ref. [14] Ruijsenaars analyzed the scattering behavior of a large class of integrable lattice systems on the discrete cone A_n (2.1) by means of stationary phase techniques. A fundamental property of the lattice systems in question is that these are diagonalized by multivariate orthogonal polynomials associated with a factorized orthogonality measure. By comparing the explicit product formula for the orthogonality measure $\Delta(\xi)$ (6.6) with the corresponding formulas in [14, Sec. 1], it is manifest that the Hall–Littlewood functions fall within the class of polynomials considered by Ruijsenaars. The upshot is that we can determine the scattering behavior of the restrictions of the Hamiltonians $H_{q,1}, \dots, H_{q,n}$ (6.9) to the n -particle subspace $\ell^2(A_n, \delta_n)$ upon describing how Ruijsenaars’ general results specialize to the present setting of the infinite q -boson model. To this end it is convenient to pass to uniform Lebesgue measures by incorporating orthogonality densities into the wave functions via the following gauge transformation:

$$\begin{aligned} \Psi_\xi(\lambda) &:= i^{n(n-1)/2} \Delta(\xi)^{1/2} \delta_n(\lambda)^{1/2} \phi_\xi(\lambda) \\ &= \delta_n(\lambda)^{1/2} \sum_{\sigma \in S_n} \text{sign}(\sigma) \hat{S}_\sigma(\xi)^{1/2} e^{i(\rho+\lambda) \cdot \xi_\sigma} \end{aligned} \tag{7.1a}$$

($\xi \in A$ (5.1)), where $\rho := \frac{1}{2}(n-1, n-3, n-5, \dots, 3-n, 1-n)$ and

$$\hat{S}_\sigma(\xi) := \prod_{\substack{1 \leq j < k \leq n \\ \sigma_j^{-1} < \sigma_k^{-1}}} s(\xi_k - \xi_j) \prod_{\substack{1 \leq j < k \leq n \\ \sigma_j^{-1} > \sigma_k^{-1}}} \overline{s(\xi_k - \xi_j)}, \tag{7.1b}$$

with $\sigma_j^{-1} := (\sigma^{-1})_j$ for $j = 1, \dots, n$ and

$$s(x)^{1/2} := \frac{1 - qe^{ix}}{|1 - qe^{ix}|}, \quad \text{so} \quad s(x) = \frac{1 - qe^{ix}}{1 - qe^{-ix}}. \tag{7.1c}$$

The wave functions in question diagonalize the commuting self-adjoint difference operators

$$\tilde{H}_{q,r} := \delta_n^{1/2} H_{q,r} \delta_n^{-1/2}, \quad r = 1, \dots, n, \tag{7.2}$$

in $\ell^2(A_n)$, viz.

$$\tilde{H}_{q,r} = \tilde{F}_q^{-1} \circ \hat{E}_r \circ \tilde{F}_q, \quad r = 1, \dots, n \tag{7.3}$$

(cf. Eq. (6.11)), where $\tilde{\mathbf{F}}_q : \ell^2(A_n) \rightarrow L^2(A, d\xi)$ denotes the Hilbert space isomorphism defined by

$$(\tilde{\mathbf{F}}_q f)(\xi) := \sum_{\lambda \in A_n} f(\lambda) \overline{\Psi_\xi(\lambda)} \quad (f \in \ell^2(A_n)) \tag{7.4a}$$

with

$$(\tilde{\mathbf{F}}_q^{-1} \hat{f})(\lambda) = \frac{1}{(2\pi)^n} \int_A \hat{f}(\xi) \Psi_\xi(\lambda) d\xi \quad (\hat{f} \in L^2(A, d\xi)) \tag{7.4b}$$

(cf. Eqs. (6.8a), (6.8b)). The action of $\tilde{\mathbf{H}}_{q,r}$ (7.2) on $f \in \ell^2(A_n)$ reads explicitly

$$\begin{aligned} (\tilde{\mathbf{H}}_{q,r} f)(\lambda) = & \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \lambda + e_J \in A_n}} V_{J,\lambda}^{1/2} V_{J^c, \lambda + e_J}^{1/2} f(\lambda + e_J) \\ & + \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \lambda - e_J \in A_n}} V_{J^c, \lambda}^{1/2} V_{J, \lambda - e_J}^{1/2} f(\lambda - e_J). \end{aligned} \tag{7.5}$$

For $1 \leq r \leq n$, let A_r be an open dense domain in A (5.1) on which the gradient vector $\nabla \varepsilon_r$ is regular with respect to the permutation-action of S_n on its components:

$$A_r := \{\xi \in A \mid \partial_j \varepsilon_r \neq \partial_k \varepsilon_r, \forall 1 \leq j < k \leq n\} \tag{7.6}$$

(with ε_r taken from Eq. (6.10b)). For any $\xi \in A_r$, there exists then a unique permutation $\sigma_\xi \in S_n$ reordering the components of $\nabla \varepsilon_r(\xi)$ in strictly decreasing order, i.e. $\sigma_\xi(\nabla \varepsilon_r(\xi)) \in \mathbb{R}_>^n := \{x \in \mathbb{R}^n \mid x_1 > x_2 > \dots > x_n\}$. Clearly the assignment $\xi \rightarrow \sigma_\xi$ is constant on the connected components of A_r by the continuity of $\nabla \varepsilon_r(\xi)$. Let $\hat{\mathcal{S}}_r$ now denote the following unitary operator on $L^2(A, d\xi)$ —the *scattering matrix*—defined via its restriction to the dense subspace of smooth test functions with compact support inside A_r :

$$(\hat{\mathcal{S}}_r \hat{f})(\xi) = \hat{\mathcal{S}}_{\sigma_\xi}(\xi) \hat{f}(\xi) \quad (\hat{f} \in C_0^\infty(A_r)). \tag{7.7}$$

The scattering matrix (7.7) allows us to compare the large-time asymptotics of the dynamics

$$(e^{it\tilde{\mathbf{H}}_{q,r}} f)(\lambda) = \frac{1}{(2\pi)^n} \int_A e^{it\varepsilon_r(\xi)} \hat{f}(\xi) \Psi_\xi(\lambda) d\xi \quad (\hat{f} = \tilde{\mathbf{F}}_q f) \tag{7.8}$$

of the higher q -boson Hamiltonian $\tilde{\mathbf{H}}_{q,r}$ (7.2) with that of the corresponding $q \rightarrow 0$ limiting Hamiltonian $\tilde{\mathbf{H}}_{0,r}$:

$$(\tilde{H}_{0,r}f)(\lambda) = \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \lambda + e_J \in \Lambda_n}} f(\lambda + e_J) + \sum_{\substack{J \subset \{1, \dots, n\}, |J|=r \\ \lambda - e_J \in \Lambda_n}} f(\lambda - e_J) \tag{7.9}$$

for the phase model of impenetrable bosons ($f \in \ell^2(\Lambda_n)$). Specifically, as a very special case of [14, Thm. 3.3] we arrive at the following explicit formulas for the corresponding wave- and scattering operators.

Theorem 7.1 (*Wave- and scattering operators*). *The operator limits*

$$\Omega_r^\pm := s - \lim_{t \rightarrow \pm\infty} e^{it\tilde{H}_{q,r}} e^{-it\tilde{H}_{0,r}} \tag{7.10a}$$

converge in the strong $\ell^2(\Lambda_n)$ -norm topology and the corresponding wave operators Ω_r^\pm are given by unitary operators in $\ell^2(\Lambda_n)$ of the form

$$\Omega_r^\pm = \tilde{F}_q^{-1} \circ \hat{S}_r^{\mp 1/2} \circ \tilde{F}_0. \tag{7.10b}$$

Consequently, the scattering operator relating the large-times asymptotics of the q -boson dynamics $e^{it\tilde{H}_{q,r}}$ for $t \rightarrow -\infty$ and $t \rightarrow +\infty$ is given by the unitary operator

$$S_r := (\Omega_r^+)^{-1} \Omega_r^- = \tilde{F}_0^{-1} \circ \hat{S}_r \circ \tilde{F}_0 \tag{7.10c}$$

($r = 1, \dots, n$).

It is instructive to outline briefly how Ruijsenaars’ proof in [14] simplifies in our particular situation. For this purpose, we associate to any $\hat{f} \in C_0^\infty(A_r)$ a ($q = 0$) boson wave packet $f^{(0)}(t)$ and q -boson wave packets $f_\pm(t)$ in $\ell^2(\Lambda_n)$ of the form

$$\begin{aligned} f^{(0)}(t) &:= \tilde{F}_0^{-1} (e^{-it\hat{E}_r} \hat{f}), \\ f_\pm(t) &:= \tilde{F}_q^{-1} (e^{-it\hat{E}_r} \hat{S}_r^{\pm 1/2} \hat{f}). \end{aligned} \tag{7.11}$$

Theorem 7.1 is now immediate from the following proposition.

Proposition 7.2 (*Asymptotic equivalence*). *For all $K > 0$ one has that*

$$\|f_\pm(t) - f^{(0)}(t)\| = O(|t|^{-K}) \quad \text{as } t \rightarrow \pm\infty \tag{7.12}$$

(where $\|\cdot\|$ refers to the ℓ^2 -norm in $\ell^2(\Lambda_n)$).

To infer the above proposition, one may assume without loss of generality that the compact support of smooth test function \hat{f} is contained inside a connected component of A_r . We then write $\hat{\sigma} \in S_n$ for the unique (ξ -independent) permutation ordering the elements of $\nabla_{\varepsilon_r}(\xi)$ in strictly decreasing order for all ξ in the support of \hat{f} . Let

$V_{\text{class}} \subset \mathbb{R}^n$ be an open bounded neighborhood of the compact range of classical wave-packet velocities $\text{Ran}(\nabla \varepsilon_r) := \{\nabla \varepsilon_r(\xi) \mid \xi \in \text{Supp}(\hat{f})\}$ staying away from the boundary of the chamber $\hat{\sigma}^{-1}(\mathbb{R}_{>}^n)$. The *classical wave packet*, finitely supported on the following t -dependent region of Λ_n :

$$\Lambda_n^{\text{clas}}(t) := \begin{cases} \{\rho + \lambda \in t\hat{\sigma}(V_{\text{clas}})\} & \text{for } t > 0, \\ \{\rho + \lambda \in t\sigma_0\hat{\sigma}(V_{\text{clas}})\} & \text{for } t < 0, \end{cases} \tag{7.13}$$

is defined as

$$f_\lambda^{\text{clas}}(t) := \begin{cases} \frac{\text{sign}(\hat{\sigma})}{(2\pi)^n} \int_A e^{i(\rho+\lambda)\cdot\xi - it\varepsilon_r(\xi)} \hat{f}(\xi) \, d\xi & \text{for } \lambda \in \Lambda_n^{\text{clas}}(t) \text{ and } t > 0, \\ \frac{\text{sign}(\hat{\sigma})}{(-2\pi)^n} \int_A e^{i(\rho+\lambda)\cdot\xi_{\sigma_0} - it\varepsilon_r(\xi)} \hat{f}(\xi) \, d\xi & \text{for } \lambda \in \Lambda_n^{\text{clas}}(t) \text{ and } t < 0, \\ 0 & \text{otherwise.} \end{cases} \tag{7.14}$$

Here σ_0 refers to the order reversing permutation for which $\sigma_j = n + 1 - j$, $j = 1, \dots, n$.

With the aid of the following stationary phase estimate from [12, pp. 38–39]: for any $K > 0$ there exists a constant $C_K > 0$ such that

$$\left| \int_{\mathbf{A}} e^{ix\cdot\xi - it\varepsilon_r(\xi)} \hat{f}(\xi) \, d\xi \right| \leq \frac{C_K}{(1 + |x| + |t|)^K} \tag{7.15}$$

for all $x \in \mathbb{R}^n$ and $t \in \mathbb{R}$ such that $x \notin tV_{\text{clas}}$, it is now not difficult to deduce that

$$\begin{aligned} \|f^{(0)}(t) - f^{\text{clas}}(t)\| &= O(|t|^{-K}) \quad \text{as } t \rightarrow \pm\infty, \\ \|f_\pm(t) - f^{\text{clas}}(t)\| &= O(|t|^{-K}) \quad \text{as } t \rightarrow \pm\infty, \end{aligned} \tag{7.16}$$

whence the asymptotic equivalence in Proposition 7.2 follows.

Acknowledgments

It is a pleasure to thank Edwin Langmann and Christian Korff for helpful conversations and comments. The referees’ constructive remarks are also greatly appreciated.

References

[1] N.M. Bogoliubov, A.G. Izergin, A.N. Kitanine, Correlation functions for a strongly correlated boson system, *Nuclear Phys. B* 516 (1998) 501–528.
 [2] A. Borodin, I. Corwin, L. Petrov, T. Sasamoto, Spectral theory for the q -Boson particle system, arXiv:1308.3475.
 [3] J.F. van Diejen, Diagonalization of an integrable discretization of the repulsive delta Bose gas on the circle, *Comm. Math. Phys.* 267 (2006) 451–476.
 [4] J.F. van Diejen, E. Emsiz, Unitary representations of affine Hecke algebras related to Macdonald spherical functions, *J. Algebra* 354 (2012) 180–210.

- [5] O. Foda, M. Wheeler, Hall–Littlewood plane partitions and KP, *Int. Math. Res. Not. IMRN* 2009 (2009) 2597–2619.
- [6] N.H. Jing, Vertex operators and Hall–Littlewood symmetric functions, *Adv. Math.* 87 (1991) 226–248.
- [7] A. Klimyk, K. Schmüdgen, *Quantum Groups and Their Representations*, Springer-Verlag, Berlin, 1997.
- [8] C. Korff, Cylindric versions of specialised Macdonald functions and a deformed Verlinde algebra, *Comm. Math. Phys.* 318 (2013) 173–246.
- [9] E.H. Lieb, W. Liniger, Exact analysis of an interacting Bose gas. I. The general solution and the ground state, *Phys. Rev. (2)* 130 (1963) 1605–1616.
- [10] I.G. Macdonald, *Symmetric Functions and Hall Polynomials*, second edition, Clarendon Press, Oxford, 1995.
- [11] I.G. Macdonald, Orthogonal polynomials associated with root systems, *Sém. Lothar. Combin.* 45 (2000/2001), Art. B45a, 40 pp.
- [12] M. Reed, B. Simon, *Methods of Modern Mathematical Physics. III. Scattering Theory*, Academic Press, New York–London, 1979.
- [13] S.N.M. Ruijsenaars, Complete integrability of relativistic Calogero–Moser systems and elliptic function identities, *Comm. Math. Phys.* 110 (1987) 191–213.
- [14] S.N.M. Ruijsenaars, Factorized weight functions vs. factorized scattering, *Comm. Math. Phys.* 228 (2002) 467–494.
- [15] T. Sasamoto, M. Wadati, Exact results for one-dimensional totally asymmetric diffusion models, *J. Phys. A* 31 (1998) 6057–6071.
- [16] N.V. Tsilevich, The quantum inverse scattering method for the q -boson model and symmetric functions, *Funct. Anal. Appl.* 40 (2006) 207–217.